

DYNAMICS OF THE NARROW-LINE REGION IN THE SEYFERT 2 GALAXY NGC 1068¹

V. DAS², D.M. CRENSHAW³, S.B. KRAEMER⁴

Accepted for publication in the Astrophysical Journal

ABSTRACT

We present dynamical models based on a study of high-resolution long-slit spectra of the narrow-line region (NLR) in NGC 1068 obtained with the Space Telescope Imaging Spectrograph (STIS) aboard *The Hubble Space Telescope* (*HST*). The dynamical models consider the radiative force due to the active galactic nucleus (AGN), gravitational forces from the supermassive black hole (SMBH), nuclear stellar cluster, and galactic bulge, and a drag force due to the NLR clouds interacting with a hot ambient medium. The derived velocity profile of the NLR gas is compared to that obtained from our previous kinematic models of the NLR using a simple biconical geometry for the outflowing NLR clouds. The results show that the acceleration profile due to radiative line driving is too steep to fit the data and that gravitational forces alone cannot slow the clouds down, but with drag forces included, the clouds can slow down to the systemic velocity over the range 100–400 pc, as observed. However, we are not able to match the gradual acceleration of the NLR clouds from ~ 0 to ~ 100 pc, indicating the need for additional dynamical studies.

Subject headings: galaxies: kinematics and dynamics – galaxies: individual
(NGC 1068) galaxies: Seyfert – AGN: [OIII] emission lines – ultraviolet: galaxies

1. INTRODUCTION

Recent studies of the kinematics of the NLRs of Seyfert galaxies have taken advantage of the high resolution of *HST* to map the velocities of these regions. (Evans et al. 1993; Macchetto et al. 1994; Hutchings et al. 1998; Nelson et al. 2000; Crenshaw et al. 2000; Crenshaw & Kraemer 2000b; Ruiz et al. 2001; Cecil et al. 2002; Ruiz et al. 2005; Das et al. 2005, 2006, and references therein). Although there have been a number of papers on the dynamical aspects of the NLR, most of these studies relied on ground-based data limited to spatial resolutions of ≥ 50 pc for even the most nearby AGN (see Kaiser et al. 2000, and references therein). These studies relied primarily upon spatially integrated line profiles to understand the dynamics of the NLR (Schulz 1990; Veilleux 1991), but the problem arose that the emission-line profiles of the NLR can be explained by many different types of dynamical models, such as infall, rotation and outflows (Capriotti et al. 1980, 1981; Vrtilek 1983; Krolik & Vrtilek 1984; Vrtilek 1985).

With the launch of *HST* and its high angular resolution ($\sim 0''.1$), the NLRs of Seyfert galaxies have received considerable attention. With the limited long-slit capability of the faint object camera (FOC), and later the expanded capability of STIS, detailed constraints on the kinematics of the NLRs in Seyfert and other galaxies became possible. In turn, these kinematic studies provided

good diagnostics upon which dynamical analyses can be based.

1.1. Previous Kinematics Studies of the NLR of Seyfert Galaxies

The structure of the NLR resembles a bicone as expected from a simple unified model of Seyfert galaxies, due to collimation by a thick torus (Antonucci & Miller 1985). Both Schulz (1990) and Evans et al. (1993) have modeled the NLR of NGC 4151 and found it to be consistent with a biconical geometry. In an *HST* study done on a sample of Seyfert 1 and 2 galaxies, Schmitt & Kinney (1996) compared both Seyfert types to study their NLR morphologies, and found triangular structures in most of their Seyfert 2s, and circular structures in most of their Seyfert 1s, consistent with the unified model and biconical structure for the NLR (see also Schmitt et al. 2003a,b). Veilleux et al. (2001) modeled the inner regions of the Seyfert galaxy NGC 2992, and found that the ionized gas can be fitted with a biconical structure.

We have recently completed a study of the kinematics of the NLRs in two Seyfert galaxies (Das et al. 2005, hereafter Paper I, and Das et al. 2006, hereafter Paper II). Since our dynamical work in this paper is a direct extension of the works of those two papers, we summarize our results below. In Paper I, kinematic models were developed to match the emission-line velocities from high-resolution STIS spectra within ~ 400 pc of the central black hole of the Seyfert 1 galaxy NGC 4151. The NLR gas showed strong evidence of acceleration from ~ 0 pc out to ~ 100 pc, then showed deceleration back to systemic velocity at ~ 400 pc with velocity roughly proportional to distance in each case. The maximum velocity of the outflowing gas at the turnover point (96 pc) was ~ 800 km s⁻¹ relative to the black hole. Based on our kinematic model, the NLR could be represented by a bicone, with inner and outer half opening angles of 15 and 33° respectively, and inclination of $\sim 45^\circ$ with respect to the plane of the sky, consistent with previous kine-

¹ Based on observations made with the NASA/ESA Hubble Space Telescope. STIS is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555.

² Indian Institute of Astrophysics, 2nd Block, Koramangala, Bangalore-560034, alvin@iiap.res.in

³ Department of Physics and Astronomy, Georgia State University, Astronomy Offices, One Park Place South SE, Suite 700, Atlanta, GA 30303, crenshaw@chara.gsu.edu

⁴ Department of Physics, Catholic University of America, 200 Hannan Hall, Washington DC, MD 20064, stiskraemer@yancey.gsfc.nasa.gov.

matic work done on NGC 4151 with different slit positions (Crenshaw et al. 2000). Some of the fainter NLR clouds showed evidence of backflow at the point where the clouds turnover in their velocities. The radio jet was found to have little effect on the kinematics of the NLR clouds, however there was some evidence of radial velocity splitting of the fainter NLR clouds near bright knots in the radio jet. The brighter clouds were not accelerated by the jet.

In Paper II, we developed a similar model for the Seyfert 2 galaxy NGC 1068 again with high-resolution spectra taken with the STIS aboard *HST*. With seven parallel slit positions covering the entire NLR, we extracted radial velocity profiles and matched them with our newly developed 3-D biconical models. Our kinematical models showed that the NLR gas accelerated out to ~ 140 pc (the turnover point), then decelerated back to systemic velocity at a distance of ~ 400 pc from the central black hole. The maximum velocity of outflow of the gas was ~ 2000 km s $^{-1}$, with respect to the black hole, and the model predicted an inner and outer half opening angles of the bicone of 20° and 40° respectively, and an inclination of 5° out of the plane of the sky. We used high resolution radio maps of the NLR obtained from Gallimore et al. (2004) to search for jet/cloud interactions. Evidence showed that, similar to NGC 4151, the fainter NLR clouds were split near bright knots in the radio jets, whereas the brighter NLR clouds remained unaffected by the jet.

Other Seyfert galaxies show similar flow patterns to NGC 4151 and NGC 1068, such as the Seyfert 2 galaxy Mrk 3 (Ruiz et al. 2001, 2005). The radial velocities of these galaxies were matched with a common kinematic model with little variation in the parameter space, which begs the question, what are the physical processes involved that would cause such a similar flow pattern in both types of Seyfert galaxies? It is this question that motivated us to carry out this dynamical study.

1.2. Previous Dynamic Studies of the NLR of Seyfert Galaxies

In the first truly dynamical model of the NLR based on *HST* kinematics, Everett & Murray (2006) attempted to fit the NLR velocities in NGC 4151, based on measurements done by Das et al. (2005). They tested an isothermal Parker wind model which assumes thermally expanding winds (Parker 1965). By assuming a spherical cloud geometry, they let the Parker wind drag along the embedded NLR clouds. As the Parker wind accelerates by thermal expansion and slowly loses heat by adiabatic cooling, the clouds are also accelerated to high speeds. They then let the Parker wind run into a low density ambient medium to slow the velocity of the wind, and hence the clouds, to the systemic velocity. The model explains the velocity profile of the NLR of NGC 4151, but suffers from the fact that an isothermal wind cannot be sustained out to large distances. Also, the mass profile of the SMBH plus galaxy, which determines the temperature profile for their model, is not exactly known for NGC 4151. Hence while their model was not successful on physical grounds, it is worth noting its relative success in matching the NLR kinematics.

In this paper, the main question we want to address is: how can we constrain the dynamics of the NLR in Seyfert

galaxies with the detailed knowledge that we have gained from our kinematic studies? Here, we concentrate on the dynamics of the NLR in NGC 1068, since it has the best constraints. We will start with a simple construction of the enclosed mass function based on data from previous studies and eventually formulate a radiation pressure-gravity tug-of-war on the NLR clouds. The questions we will attempt to answer include the following: 1) If the NLR gas is in outflow, is radiation pressure really the best driving mechanism? 2) If the NLR gas is turning over its velocity and decelerating back to systemic, is gravity responsible for stopping the gas? 3) Can we fit the velocity profile of the data with a simple radiation-gravity law, or do we need to include another force (such as drag)?

Our analysis applies to NGC 1068 in particular, but has relevance to all Seyferts in general that show signs of gas outflow and subsequent deceleration (Ruiz et al. 2005). To test whether gravity is playing any role in stopping and turning back the gas velocity, we will construct a mass profile within $\sim 10,000$ pc from the nucleus of NGC 1068. With the mass profile in hand, we will test whether the gas kinematics are dominated by rotation. Such a test might prove fruitful for rotation in NGC 4151, which shows redshifts northeast and blueshifts southwest of the nucleus, but from a geometrical point of view, rotation of the gas will prove difficult to match the velocity field of NGC 1068, which shows blueshifts and redshifts on each side of the nucleus. Next we plot the escape velocity with distance to see if the gas should escape or not, given the velocities seen in the data, and whether or not the kinematics of the NLR can be dominated by gravitational infall. Then we concentrate on outflow assuming spherical symmetry and pure radial motion. First we determine if the deceleration of the gas can be attributed to the enclosed mass, regardless of the outward accelerating force. We then apply radiative line driving plus gravitational forces and compare the results to the observed velocity law of NGC 1068 derived from the kinematic models. Finally, we introduce a drag force due to an ambient medium on the NLR clouds, in addition to radiation pressure and gravity, to determine if it can improve the fit to either the accelerating or decelerating portions of the observed velocity curve.

2. BUILDING THE MASS PROFILE

In building a model of the enclosed mass as a function of distance from the central SMBH, we incorporate various subsystems into our mass profile. These include contributions from the SMBH, the nuclear stellar cluster, and the bulge. For all these systems, we assume spherical symmetry for simplicity. We have assumed that the stellar cluster and bulge extend all the way inward to the SMBH, which may overestimate the mass close in. The size of the stellar cluster was estimated to be ~ 140 pc based on a study by Thatte et al. (1997) and the bulge was assumed to extend up to $10,000$ pc. Mass contribution from the galactic disk of NGC 1068 was neglected because the galactic potential is dominated by the large bulge to at least 1000 pc, which is well beyond the extent of the NLR.

2.1. The Black Hole Mass

NGC 1068 is one of only a few AGN that shows an edge-on disk of H₂O maser emission close to the SMBH. The disk shows the signature of a rotational velocity curve, which can be used to determine the mass of the SMBH. Greenhill & Gwinn (1997) estimated the mass of NGC 1068 to be $1.5 \times 10^7 M_\odot$ within 0.65 pc, based on the velocity field of the H₂O maser emission observed with the VLBA and the VLA. We will therefore use this estimate for the mass of the SMBH.

2.2. The Bulge Mass

Since we are going out to $\sim 10,000$ pc, which encloses the NLR and the ENLR clouds, we need an accurate assessment of the total mass within this region. At small distances (≤ 1 pc), the SMBH is dominant, although in the case of NGC 1068, a concentrated stellar cluster is also providing substantial gravity close in. For an estimate of the bulge mass in NGC 1068, we rely on the work of Häring & Rix (2004). They found a tight correlation between black-hole mass and bulge mass for a sample of 30 galaxies, including NGC 1068. They determined the bulge mass by modeling the bulge with the Jeans equation in spherical form. They assumed the bulge to be isotropic and spherically symmetric, which might lead to an overestimation of the bulge mass; however they also neglect any contribution from dark matter, which would tend to underestimate the bulge mass. Their value for the bulge mass in NGC 1068 is $2.3 \times 10^{10} M_\odot$ within a radius $r = 3R_e$ (3 effective radii), where $R_e = 3.1 \pm 0.8$ kpc, a value taken from the surface brightness deconvolution of Marconi & Hunt (2003). The effective radius R_e is defined to be such that half of the total light from the galaxy is predicted to be contained within the isophotal ellipse that has area πR_e^2 (Binney & Merrifield 1998).

Elliptical galaxies' and bulges' surface brightnesses can be well described by the empirical formula developed by de Vaucouleurs (1948)

$$I(R) = I_e e^{-7.6692[(\frac{R}{R_e})^{\frac{1}{4}} - 1]}, \quad (1)$$

where $I_e = I(R_e)$. In the 1980s–1990s, a family of stellar density curves emerged that modeled both elliptical galaxies and bulges well. These curves are of the form

$$\rho(r) = \frac{(3-\gamma)M}{4\pi} \frac{\eta}{r^\gamma(r+\eta)^{4-\gamma}}, \quad (2)$$

where η (in pc) is a scaling radius and M is the total mass of the bulge (Dehnen 1993). The parameter γ determines different types of models, where the $\gamma = 2$ cases corresponds to previous density models by Jaffe (1983) and the $\gamma = 1$ cases to models by Hernquist (1990). These models, when integrated over a spherical volume, yield the following enclosed mass profile:

$$\begin{aligned} M(r) &= \int_0^r 4\pi t^2 \rho(t) dt = \frac{4\pi(3-\gamma)M\eta}{4\pi} \int \frac{t^2 dt}{t^\gamma(t+\eta)^{4-\gamma}} \\ &= M \left(\frac{r}{r+\eta} \right)^{3-\gamma}. \end{aligned} \quad (3)$$

For the special case when $\gamma = 1.5$, the density profile of Equation 2 yields a surface density distribution that closely matches the de Vaucouleurs surface brightness profile of Equation 1 to within 15% over nearly 4 decades in radius (Dehnen 1993). Therefore we adopt

the following form of the mass function as the bulge profile

$$M(r) = M \left(\frac{r}{r+\eta} \right)^{1.5}. \quad (4)$$

We find a suitable value for η in Equation 4 by first defining the 'half-mass radius' $r_{\frac{1}{2}}$:

$$\frac{1}{2}M = M \left(\frac{r_{\frac{1}{2}}}{r_{\frac{1}{2}}+\eta} \right)^{\frac{3}{2}}, \quad (5)$$

which yields the following relation for η

$$\eta = r_{\frac{1}{2}}(2^{\frac{2}{3}} - 1). \quad (6)$$

Dehnen (1993) has found a simple approximation for $\frac{R_e}{r_{\frac{1}{2}}}$ that depends only slightly on γ . For a $\gamma \leq \frac{5}{2}$, he found that

$$\frac{R_e}{r_{\frac{1}{2}}} \approx (0.7549 - 0.00439\gamma + 0.00322\gamma^2 - 0.00182\gamma^3). \quad (7)$$

Therefore we can find a value for η by using $\gamma = 1.5$, R_e given above, and Equation 6. We find that $\eta \sim 2400$ pc. We also know that $M(3R_e) = 2.3 \times 10^{10} M_\odot$, so that

$$\begin{aligned} M &= 2.3 \times 10^{10} / \left[\frac{3 \times 3.1 \times 10^3}{3 \times 3.1 \times 10^3 + 2400} \right]^{1.5} \\ &= 3.2 \times 10^{10} M_\odot. \end{aligned} \quad (8)$$

The bulge mass distribution can finally be written as

$$M(r) = 3.2 \times 10^{10} \left(\frac{r_{pc}}{r_{pc} + 2400} \right)^{1.5} M_\odot. \quad (9)$$

2.3. The Nuclear Stellar Cluster Mass

It is known that NGC 1068 has a compact nuclear stellar cluster ~ 140 pc in radius (Thatte et al. 1997; Crenshaw & Kraemer 2000a), which contributes significant mass to the total mass profile of the NLR of NGC 1068. Therefore this mass must be taken into account when deriving the mass profile. Thatte et al. (1997) found a mass of $6.8 \times 10^8 M_\odot$ within $1''$ of the SMBH of NGC 1068, assuming a virialized, isotropic, and isothermal distribution of the stars. They used the stellar velocity distribution (σ_*) found in Dressler (1984), a value of 143 ± 5 km s⁻¹ at $\sim 1''$, to calculate the total dynamical mass within $1''$ of the nucleus of NGC 1068.

$$M_{dyn} = \frac{2\sigma_*^2 R}{G}, \quad (10)$$

where $R = 1'' \approx 72$ pc for NGC 1068. This mass includes contributions from the stellar cluster, the nucleus, and the bulge within a radius of 72 pc. Therefore to find just the mass from the stellar cluster M_{sc} , we took out the rest of the mass contribution from the total mass.

$$M_{sc}(72pc) = M_{dyn}(72pc) - M_{smbh} - M_{bulge}(72pc) \quad (11)$$

Making the various substitutions we have

$$\begin{aligned} M_{sc}(72pc) &= 6.8 \times 10^8 - 1.5 \times 10^7 \\ &\quad - 3.2 \times 10^{10} \left(\frac{72}{72 + 2400} \right)^{1.5} \\ &= 5.1 \times 10^8 M_\odot. \end{aligned} \quad (12)$$

Based on the radial surface brightness profile presented in Figure 4 of Thatte et al. (1997), Beckert & Duschl (2004) computed a power law consistent with the form $S_*(r) \propto r^{-1}$ to fit the data. They claim that if the profile traces the stellar mass distribution, then they would expect that the mass profile would have the form $M_*(r) \propto r$, given a spherical, isothermal distribution of stars in the cluster. Using their distribution we can estimate the stellar mass function based on the condition $M_{sc,72} = 5.1 \times 10^8 M_\odot = kr$, so that $k = \frac{5.1 \times 10^8}{72} = 7.1 \times 10^6 M_\odot \text{ pc}^{-1}$. The stellar cluster mass function is therefore

$$M_{sc}(r) = 7.1 \times 10^6 r_{pc} M_\odot. \quad (13)$$

Beyond the observed extent of the stellar cluster, we assumed that $M_{sc}(r)$ is a constant. Finally the total enclosed mass function for the NLR of NGC 1068 is given by $M_{tot}(r) = M_{smbh} + M_{bulge} + M_{sc}$ or

$$M_{tot}(r) = 1.5 \times 10^7 + 7.1 \times 10^6 r_{pc} + 3.2 \times 10^{10} \left(\frac{r_{pc}}{r_{pc} + 2400} \right)^{1.5} M_\odot. \quad (14)$$

3. DYNAMICS BASED ON GRAVITY

A figure representing the total mass enclosed within r_{pc} is shown in the top panel of Figure 1. The mass profiles for each contribution, the SMBH, bulge, and cluster are also shown in the figure. Close in toward the nucleus, the nuclear stellar cluster dominates up to its entire extent, while the bulge takes over from there. The black hole mass dominates at ≤ 2 pc. The kink in the total mass curve at ~ 140 pc is because the stellar cluster was cut off abruptly at this location. We could have modeled the cluster to assume an exponential drop-off after 140 pc, but this would not have contributed much to the gravitational force exerted by the total mass.

3.1. Rotation

With the total mass profile we calculate the circular rotational velocity and plot it as a function of distance as shown in the middle panel of Figure 1. The rotation velocity only depends on the enclosed mass at radius r_{pc} , and is given by the formula below

$$V(r) = \sqrt{G \frac{M(r)}{r}}. \quad (15)$$

For demonstration purposes, we show the rotational velocities as if only each mass component was present, as well as the velocity for the total mass profile. The rotational velocity profile indicates to us that rotation cannot dominate the kinematics of the NLR of NGC 1068 because many observed data points exhibit large velocities ($\geq 1000 \text{ km s}^{-1}$), whereas the rotation curve never exceeds $\sim 220 \text{ km s}^{-1}$ at large distances (100 pc). Again the stellar cluster dominates the velocities up to ~ 300 pc.

3.2. Escape and Infall Velocity

The escape velocity at a given distance r_{pc} is calculated numerically from the formula

$$V(r) = \sqrt{\int_r^\infty 2G \frac{M(t)}{t^2} dt} \text{ km s}^{-1}, \quad (16)$$




FIG. 1.— Top: The total enclosed mass profile of NGC 1068 in solid black. The individual contributions to the total mass is also shown in this plot. They are the stellar cluster (dash-dot), the bulge (dash), and the SMBH (dot). The stellar cluster was modeled to have an extent of 140 pc. Middle: The total rotational velocity profile of the NLR of NGC 1068 in solid black color. The individual rotation components due to the various masses are also shown. Bottom: The escape velocity as a function of distance for the NLR of NGC 1068.

based on the enclosed mass function, and is plotted in the bottom panel of Figure 1. According to our kinematic model, the maximum velocity at the turnover radius $r_t = 140$ pc is 2000 km s^{-1} (Paper II); therefore Figure 1 tells us that the NLR clouds should have escaped after 140 pc, where the escape velocity is only $\sim 500 \text{ km s}^{-1}$. In the data however, clouds at 140 pc are at much higher velocity than escape velocity; yet after the turnover point, the clouds start to decrease their velocity and return to systemic. Thus, some force other than gravity is causing the clouds to decelerate at $r \geq 140$ pc. The infall velocity profile is equivalent to the escape velocity profile, except that the velocity vector is now directed inward. Therefore gravitational infall also cannot account for the faster moving clouds at 140 pc, and in general does not match the observed velocity profile (Paper II).

3.3. Gravitational Drag

To test the importance of the force of gravity alone on slowing down the outflowing NLR clouds, we give the clouds a maximum velocity at the turnover point and let gravity do the rest. In other words, we assume that there is no outward driving force after the turnover point and let the clouds coast under the force of gravity. The top panel of Figure 2 shows that with a maximum velocity $\geq 1000 \text{ km s}^{-1}$ at 140 pc, there is little deceleration with radius. However, with maximum velocities $\leq 300 \text{ km s}^{-1}$ gravity can slow the clouds down to rest, as seen in the bottom panel of Figure 2. The maximum velocities in the NLRs of some Seyfert galaxies are on the order of $\sim 400 \text{ km s}^{-1}$ (Ruiz et al. 2005), and gravitational deceleration may be important in these cases. The kinematic model of NGC 1068, however, shows maximum velocities of up to $\sim 2000 \text{ km s}^{-1}$, clearly out of the reach of gravitational deceleration. Gravity alone cannot slow down the outflowing clouds in this case and there must be some other force or forces involved.

To compound the problem, suppose we let radiation or some other force push on the gas while gravity is trying to pull it back. In this case the gravitational deceleration will be even less. However, to further explore this issue, we generate a line-driven radiation model, with gravity competing to slow the gas down. This will eventually lead us in the direction of including a drag force to complete the analysis.

4. RADIATION-GRAVITY FORMALISM

The various mechanisms to push the gas out from close to the nucleus include a radiation pressure driven wind, a thermally driven wind, or a magneto-hydrodynamic (MHD) wind (Crenshaw et al. 2003). The latter two methods are discussed in more detail in Everett (2005) and Everett & Murray (2006), and were summarized in §1.2. However, no dynamical model to date has led to a satisfactory description of the kinematics in NGC 1068 or NGC 4151. The radiation driven wind mechanism is most efficient when the momentum imparted to the gas is due to line-driving (bound-bound transitions), although bound-free and free-free electron transitions (Thomson scattering) also contribute to driving the gas out (Chelouche & Netzer 2001). The radiation force is dependent on the ionization state of the gas, with lower ionization states more efficient due to the greater availability of electrons in the bound states. If dust is mixed




FIG. 2.— The velocity slowing down with distance for an initial launch at 140 pc with 1000 km s^{-1} (top) and 400 km s^{-1} (bottom) shown in solid lines for the clouds in NGC 1068. The other mass components' effect are shown in different linestyles represented by the key above the curves. The clouds barely slow down out to 10,000 pc when launch with 1000 km s^{-1} . However, gravity was able to slow down the clouds down when they were launched with 300 km s^{-1} .

in with the NLR gas, then it will compete with the gas in absorbing ionizing photons and hence radiation pressure on the dust can become an important contributor to the velocities of the outflowing gas in the NLR (Dopita et al. 2002). If the dust grains are electrically charged, they can drag the ionized gas along to similar velocities as the dust. In this section we ignore the effects of dust, which would only increase the radiative acceleration. Thus we consider a radiation driving mechanism coupled with the effects of gravity to find the velocity profile of the NLR gas.

4.1. Building the Velocity Equation

We start with the acceleration due to radiation on a point mass,

$$a(r) = \frac{L\sigma_T\mathcal{M}}{4\pi r^2 c m_p}, \quad (17)$$

where a is the acceleration, L is the bolometric luminosity of NGC 1068, σ_T is the Thomson scattering cross section for the electron, r is the distance, c is the speed of light, m_p is the mass of the proton, and $\mathcal{M}$ is the force multiplier. As mentioned above, to really drive the gas out efficiently, we need to incorporate other sources of opacity such as bound-bound and bound-free opacity in addition to those from Thomson scattering. These additional opacities are included via the force multiplier, $\mathcal{M}$ in Equation 17, which is primarily a function of ionization parameter⁵ U for a given spectral energy distribution (Crenshaw et al. 2003, and references therein).

The acceleration due to gravity per mass is simply given by

$$a(r) = -\frac{GM(r)}{r^2}, \quad (18)$$

where $M(r)$ is the total enclosed mass within r parsecs, and G is the universal gravitational constant. Putting Equations 17 and 18 together we have

$$a(r) = \frac{L\sigma_T\mathcal{M}}{4\pi r^2 cm_p} - \frac{GM(r)}{r^2}. \quad (19)$$

Now we will have to rewrite the acceleration in terms of velocity as a function of radius, and then solve for $v(r)$.

$$a = \frac{dv}{dt} = \frac{dr}{dt} \frac{dv}{dr} = v \frac{dv}{dr}. \quad (20)$$

Therefore we can now write Equation 19 as a simple separable differential equation

$$v dv = \frac{L\sigma_T\mathcal{M}}{4\pi r^2 cm_p} dr - \frac{GM(r)}{r^2} dr. \quad (21)$$

Substituting for the constants in Equation 21 and converting to appropriate units of km s^{-1} and pc, then integrating and setting the initial velocity to zero yields the following form for $v(r)$

$$v(r) = \sqrt{\int_{r_1}^r \left[6840 L_{44} \frac{\mathcal{M}}{t^2} - 8.6 \times 10^{-3} \frac{M(t)}{t^2} \right] dt}, \quad (22)$$

where L_{44} is luminosity in units of $10^{44} \text{ ergs s}^{-1}$ and $M(r)$ is in units of $M_\odot$. The constraints on the luminosity and the force multiplier are presented in the next section.

4.2. Physical Constraints on the NLR of NGC 1068

Since NGC 1068 is a Seyfert 2 galaxy, we cannot measure its luminosity directly. From Pier et al. (1994), the total luminosity of NGC 1068 is given by

$$L_{bol} = 2.2 \times 10^{11} \left(\frac{f_{refl}}{0.01} \right)^{-1} \left(\frac{D}{22 \text{ Mpc}} \right)^2 L_\odot, \quad (23)$$

where f_{refl} is the fraction of nuclear flux observed as scattered radiation, D is the distance to NGC 1068, and $L_\odot$ is the solar luminosity. The most uncertain term in Equation 23 is f_{refl} . Pier et al. (1994) have summarized a range of values for f_{refl} that have been determined previously by several authors. The range in f_{refl} spans a few

orders of magnitude, from 0.001–0.05. They claim that the best estimate comes from Miller et al. (1991), who had determined a value for f_{refl} to be 0.015, based on observations of [OIII] and broad $H\beta$ luminosity and their ratio. Pier et al. (1994) concluded that f_{refl} is probably within a factor of a few of 0.01, hence we have adopted a value for f_{refl} of 0.015 because it is the “best” estimate and close to the average value adopted by Pier et al. (1994). We already know the distance to NGC 1068 as 14.4 Mpc (Bland-Hawthorn et al. 1997) so the bolometric luminosity of NGC 1068 is given by

$$L_{bol} = 2.2 \times 10^{11} \left(\frac{0.015}{0.01} \right)^{-1} \left(\frac{14.4}{22 \text{ Mpc}} \right)^2 L_\odot \\ = 2.4 \times 10^{44} \text{ erg s}^{-1}, \quad (24)$$

a value which could be uncertain by a factor of 0.3–15, depending on f_{refl} .

The emission lines arising from the NLR gas are best fitted with a two component photoionization model at each position, based on *HST*/STIS long-slit spectra (Kraemer & Crenshaw 2000). According to Kraemer & Crenshaw, the ionization state of the gas ranges from $U \sim 10^{-1.5}$ – $10^{-3.0}$ for the two components but seems to vary little with distance. Using U and the SED for NGC 1068, we found the force multiplier to vary from $\mathcal{M} \approx 500$ –6000 for the front face (ionized face) of the clouds, based on CLOUDY models (Ferland et al. 1998). The mass function was derived in previous sections, and $L_{44} = 2.4$, so we can now numerically solve Equation 22 for $v(r)$, assuming $\mathcal{M}$ is constant with distance. The results are presented in the next section.

4.3. Radiation and Gravity Results

Equation 22 has only two parameters that we can vary to find the velocity v as a function of distance r : the launch radius r_1 , and the force multiplier $\mathcal{M}$. We plotted $v(r)$ for various combinations of launch radii and force multipliers of the gas. The top panel of Figure 3 shows that with a launch radius of $r_1 = 1$ pc, the velocity of the gas increases with force multiplier, but quickly reaches a terminal velocity and does not slow down significantly, even out to $\sim 10,000$ pc. This is not a surprise, as previous plots had shown that the mass is not enough to slow down the high-velocity clouds. In addition, the maximum velocities for $\mathcal{M} \geq 500$ are too high compared to the observations, indicating that we must increase the launch radius r_1 .

The physical constraints on the NLR gas suggested that the force multiplier is larger than 500. However, we generated plots for $\mathcal{M}$ smaller than 500 to see whether or not gravity can slow the clouds down, if the clouds are launched from 1 pc. In the middle panel of Figure 3, we see that with force multipliers ≤ 40 , gravity does slow the clouds down, but that the maximum velocities are too low to fit the data of NGC 1068. Such models may prove useful for Seyferts with lower outflow velocities in their NLRs, such as NGC 4151 and Mkn 3 (Das et al. 2005; Ruiz et al. 2001).

We next fix the force multiplier to the lowest value of 500 (because higher values of $\mathcal{M}$ will be even more problematic to slow down the clouds) and vary the launch radius. These plots are presented in the bottom panel of Figure 3. The maximum velocity falls below the observed values at $r_1 \geq 2$ pc, and the final velocity does not

⁵ The ionization parameter U is defined as $U = \frac{\int_{\nu_0}^{\infty} \frac{L_\nu}{4\pi r^2 n_H c} d\nu}{\frac{\# \text{ of ionizing photons}}{\# \text{ H atoms}}}$ at the ionized face of the clouds, where $h\nu_0 = 13.6 \text{ eV}$ and n_H is the hydrogen number density.




FIG. 3.— Top: The velocity profiles for several force multipliers $\mathcal{M}$ with a fixed launch radius $r_1 = 1$ pc. None of the curves show a significant decrease in velocity up to 10,000 pc. Middle: The velocity profiles for several lower force multipliers with $r_1 = 1$ pc. Curves with force multipliers less than 40 turn over before reaching 10,000 pc, although their maximum velocities do not fit the data of NGC 1068. Bottom: The velocity profiles for several different launch radii (top to bottom curves $r_1 = 1, 2, 4, 8, 16, 32$ pc) all modeled with a constant force multiplier of 500. The bottom curve was able to turn over and return to systemic at ~ 1000 pc. The maximum velocity and maximum distance reached for these curves do not fit the data.

return to zero until we reach large launch radii. As the launch radius increases to ~ 32 pc, the maximum attainable velocity of the outflowing clouds decreases, and the enclosed mass is finally able to turn the velocity around. However by that time, the maximum velocity is much lower than predicted by the kinematic model. Also, the launch radius and the maximum distance of outflow become too large to fit the data, since the NLR clouds generally launch from close in and drop in radial velocity within 400 pc (Paper II). Clearly these curves cannot fit the observed NLR velocities of NGC 1068.

5. RADIATION, GRAVITY, AND DRAG FORCES

The radiation-gravity interaction on the NLR clouds of NGC 1068 fails to reproduce its velocity profile. For reasonable parameters, the velocity increases quickly close to the nucleus and mostly remains constant over large distances regardless of launch radius. The maximum outflow velocity is rather sensitive to launch radius and decreases with increasing r_1 but the clouds' velocities never turn over and decrease. Fine-tuning r_1 and $\mathcal{M}$ outside of the range of reasonable parameters can lead to a deceleration profile, but the resulting velocity profile and amplitude does not match the observed trend. Therefore we conclude that there must be additional forces at play in the NLR to account for the velocity profile that we see in the data. One such force that could explain the trend in the data is drag, whereby the clouds are slowing down in a more diffuse, hotter, and higher ionization medium (Crenshaw & Kraemer 2000b).

5.1. Deceleration due to Drag

The drag force exerted on a cloud by an ambient medium is

$$F_{\text{drag, cloud}} = \rho_{\text{med}}(v_{\text{cloud}} - v_{\text{med}})^2 A_{\text{cloud}}, \quad (25)$$

where ρ_{med} is the mass density of the ambient medium, v_{cloud} and v_{med} are the velocities of the cloud and medium respectively, and A_{cloud} represents the cross sectional area of a cloud (Everett & Murray 2006). Following Everett & Murray, we assume the clouds to be spherical with mass $m_{\text{cloud}} = \frac{4}{3}\pi R_{\text{cloud}}^3 \rho_{\text{cloud}}$, where R_{cloud} is the radius of a cloud, and ρ_{cloud} is its mass density. The acceleration on the clouds due to drag is then

$$\begin{aligned} a_{\text{drag, cloud}} &= \frac{\rho_{\text{med}}(v_{\text{cloud}} - v_{\text{med}})^2 \pi R_{\text{cloud}}^2}{\frac{4}{3}\pi R_{\text{cloud}}^3 \rho_{\text{cloud}}} \\ &= \frac{m_p n_{\text{med}}}{m_p n_{\text{cloud}} R_{\text{cloud}}} \frac{3}{4} (v_{\text{cloud}} - v_{\text{med}})^2, \end{aligned} \quad (26)$$

where n is the hydrogen number density and m_p is the mass of the proton. In our case, we assume that the velocity of the ambient medium is zero and the radius of the cloud remains constant. Therefore with $v_{\text{med}} \approx 0$ we can write

$$a_{\text{drag, cloud}} = -\frac{n_{\text{med}}}{N_{\text{H, cloud}}} \frac{3}{4} v_{\text{cloud}}^2, \quad (27)$$

where $N_{\text{H, cloud}} (\approx n_{\text{cloud}} R_{\text{cloud}})$ is the hydrogen column density of the cloud, and the minus sign represents deceleration. Combined with the radiative and gravitational acceleration of Equation 19, the total acceleration on the clouds becomes

$$a_{\text{tot}} = \frac{L_{\text{ST}} \mathcal{M}}{4\pi r^2 c m_p} - \frac{GM(r)}{r^2} - \frac{n_{\text{med}}}{N_{\text{H, cloud}}} \frac{3}{4} v_{\text{cloud}}^2, \quad (28)$$

and in differential form is

$$v dv = \frac{L_{\sigma T} \mathcal{M}}{4\pi r^2 c m_p} dr - \frac{GM(r)}{r^2} dr - \frac{n_{med}}{N_{H,cloud}} \frac{3}{4} v^2 dr. \quad (29)$$

Substituting the various constants and converting to units of pc and km s^{-1} , we have the following inseparable differential equation to solve

$$\begin{aligned} \frac{dv(r)}{dr} = & 3420 L_{44} \frac{\mathcal{M}}{v(r) r^2} - 4.3 \times 10^{-3} \frac{M(r)}{v(r) r^2} \\ & - 2.3 \times 10^{-2} \frac{n_{med}}{N_{20}} v(r), \end{aligned} \quad (30)$$

where $N_{20} = N_H / 10^{20} \text{cm}^{-2}$.

5.2. Constraints on Cloud and Medium Densities for NGC 1068

Both NGC 1068 and NGC 4151 show evidence for highly ionized gas extended throughout their NLRs, based on Chandra X-ray Observatory images (Ogle et al. 2000, 2003). Thus, we assume an ambient medium that is highly ionized, with ionization parameter $U \approx 10$. The ionization parameter, which is inversely related to the density and radius, can be written as follows

$$U \propto \frac{\int_{\nu_0}^{\infty} \frac{L_{\nu}}{h\nu} d\nu}{r^2 n_H} \quad (31)$$

Therefore if the ambient and the NLR clouds see the same ionizing luminosity (L_{ion}) at a particular distance r from the source, we can write

$$n_{med} = n_{cloud} \frac{U_{cloud}}{U_{med}}. \quad (32)$$

Kraemer & Crenshaw (2000) provided good constraints on the parameters on the right side of Equation 32. The densities of the NLR clouds are almost constant out to large distances from the nucleus with a typical value of $n_{cloud} \approx 10^4 \text{cm}^{-3}$ for clouds with $U_{cloud} \approx 10^{-3}$. If we substitute these estimates in Equation 32, we will have a typical estimate for n_{med} as follows

$$n_{med} = 10^4 \frac{10^{-3}}{10} = 1 \text{cm}^{-3}. \quad (33)$$

Kraemer & Crenshaw found column densities for the NLR clouds in the range $N_H = 10^{19} - 10^{21} \text{cm}^{-2}$, which corresponds to $N_{20} = 0.1 - 10$. Since we are interested in the ratio $\frac{n_{med}}{N_{20}}$, varying either parameter while keeping the other constant will result in the same curves. In this paper, we choose to keep N_{20} constant at the average value of 1 and vary n_{med} , because n_{med} is the most unknown quantity. We already know that $\mathcal{M}$ can take values from 500–6000, so Equation 30 can now be solved numerically for $v(r)$ with various values of the launch radius (r_1), force multiplier ($\mathcal{M}$), and the ratio $\frac{n_{med}}{N_{20}}$. We used Mathematica v5.2, which employs the most efficient choices among various flavors of Runge-Kutta algorithms, to solve for $v(r)$. The results are presented in the next section.

5.3. Radiation, Gravity and Drag Results

The top panel of Figure 4 represents several plots with a force multiplier of 500, a launch radius of 1 pc, a column density of $N_{20} = 1$, and various densities of the




FIG. 4.— Top: The velocity profiles for force multiplier 500, a launch radius of 1 pc, and varying medium densities n_{med} shown by the numbers. Bottom: The velocity profiles for a force multiplier of 6000, a launch radius of 20 pc, and varying medium densities similar to the top panel.

ambient medium. The figure shows that when launched at 1 pc, the gas accelerates to a maximum velocity inside 10 pc, then slows down again. The velocity slows down faster with increasing drag forces, as measured by increasing densities of the ambient medium. However the point of maximum velocity is too close in to match the data, which has a turnover velocity at ~ 140 pc. The curve with medium density $n_{med} = 0.33$ shows gradual deceleration out to ~ 500 pc, but its maximum velocity is slightly too high. Any other curve has either too high velocity at turnover or the velocity drops too quickly. Note that the velocity of the data we are trying to match has $\sim 2000 \text{km s}^{-1}$ at a turnover of ~ 140 pc for NGC 1068. With force multipliers higher than 500, the same trend as in Figure 4 is seen in the velocity model, except that the velocities are much higher ($\geq 6000 \text{km s}^{-1}$). Again, the curve that seems best to match the deceleration part in the data is with medium density $n_{med} = 0.333$. The rest of the curves decelerate too quickly or too slowly. The major problem with all of these curves is that the velocity turnover point is much closer to the nucleus than

TABLE 1
PARAMETERS USED TO GENERATE THE DRAG MODEL SHOWN IN
FIGURE 5.

$z_{\max}$ (pc)	θ_{inner} (deg)	θ_{outer} (deg)	i_{axis} (deg)	PA_{axis} (deg)	v_{law}
450	10	40	5	57.8	Equation 30

$z_{\max}$ is the distance from the center to one end of the bicone, along the z-axis.

θ_{inner} is the inner opening angle of the bicone.

θ_{outer} is the outer opening angle of the bicone.

i_{axis} is the inclination of the bicone axis out of the plane of the sky, with positive inclination implied by the north bicone closer toward the observer.

PA_{axis} is the position angle of the bicone axis in the plane of the sky.

v_{law} is the velocity law used in mapping the velocity field onto the bicone.

140 pc.

Another way to decrease the overall velocity is to increase the launch radius, and tweak the force multiplier and ambient density. We tried a launch radius of 10 pc with force multiplier ranging from 500–6000. The maximum velocity drops as expected, reaching $\sim 700 \text{ km s}^{-1}$ for a force multiplier of 500 and climbs to $\sim 2500 \text{ km s}^{-1}$ when we increase the force multiplier to 6000. We noticed that the turning point increases by a factor of 5 as we increase the launch radius from 1 to 10 pc. However the turnover point is still too low to fit the data well. In order to have a good fit, we first need to maximize the launch radius and then tweak the force multiplier and medium density to get the closest match to the data as possible.

The resolution in our data is ~ 10 pc and we should not launch much beyond this distance since we see clouds close to the SMBH with near zero velocity. Therefore, in the bottom panel of Figure 4, we present a plot with a launch radius of $r_1 = 20$ pc, a force multiplier of 6000, and a column density of $N_{20} = 1$. The maximum velocity reached is $\sim 1700 \text{ km s}^{-1}$ with $n_{\text{med}} = 10$. The best curve to represent the data seems to be the one with $n_{\text{med}} = 0.33$, whose maximum velocity is $\sim 1500 \text{ km s}^{-1}$, although the turnover point at ~ 60 pc is still too low according to our kinematic model. Furthermore, we have had to tweak the launch radius and force multiplier to very specific values, such that only a narrow range of the observed values gives a reasonably decent fit. However, to directly test this velocity profile, we generate biconical models similar to our previous kinematic models (Paper II), to determine whether a more dynamical velocity law, rather than the simple linear law, can reasonably match the data.

5.4. Dynamical Fit

We applied our kinematic models of Paper II with our dynamical velocity law to the data. Previously we had used a simple kinematic velocity law, which is based on the relation $v = kr$. Now instead we substitute the velocity relation based on Equation 30, which represented a more physical situation. A model with the new velocity law of Equation 30 is presented in Figure 5, using the input parameters from Table 1, and using the best-fit curve from the bottom panel of Figure 4, with $N_{20} = 1$ and $n_{\text{med}} = 0.33$. Our biconical model was constructed




FIG. 5.— The model of slit 4 generated with input parameters from Table 1, and the best-fit velocity profile from the bottom panel of Figure 4 with $n_{\text{med}} = 0.33$. Clearly this model is a poor match to the data.

with an inner and outer half opening angle of 10° and 40° respectively, and inclination of 5° , and position angle of 57.8° , and a half-size of 450 pc. Points on the bicone were assigned velocities according to Equation 30. The center slit (slit 4, see Paper II) is used here for comparison. We extracted the velocities from the model at a position equivalent to slit 4 and those are shown in the shaded regions in Figure 5. The data from slit 4 are shown in small triangles.

The model represents a poor fit to the data from slit 4 (the center slit) of NGC 1068. This was expected from looking at the previous velocity plots, as the turnover point was too low, the launch radius was a bit large, and the velocity profile did not resemble our kinematically derived linear profiles. We have varied the input parameters from Table 1, but this makes very little improvement. The turnover point, starting distance, and maximum velocity of the bicone cannot be varied without changing the drag parameters, as these are implicitly defined in Equation 30. The thickness of the bicone can be increased to accommodate more data points, but the bicone will show lots of unnecessary shaded regions. Changing the maximum extent of the bicone $z_{\max}$ will have absolutely no effect on the shaded region, except interrupting the shaded regions before or continuing them beyond $\pm 6''$. That leaves us with only two parameters to vary, the inclination of the bicone axis, and its position angle in the sky, and varying these two alone did not fix the model.

6. CONCLUSIONS

With radiation pressure driving the NLR clouds, their velocities will accelerate very quickly, within a few parsecs of the nucleus, assuming the clouds are indeed launched close in. With the introduction of the drag forces, the overall velocities are lowered, but even so the velocities reach maximum too quickly. The data suggest that the clouds are gradually increasing their velocities to a maximum at about 140 pc. This gradient in the ve-

locity cannot be simply accounted for by radiative forces driving the clouds. It seems therefore, that radiation pressure may not be the only driving mechanism for the NLR clouds, or that other forces are involved to steer the clouds to that particular gradient. The velocity profile shown in the data resembles one with a linear or ‘Hubble flow’ law. The inclusion of a drag force only serves to reduce the overall high velocities due to radiation driving, but the maximum velocities are reached very close to the nucleus, too close to match the data effectively.

Gravitational forces alone cannot stop the fast moving clouds observed in the NLR of NGC 1068. With velocities as high as 1500 km s^{-1} at 140 pc, the clouds should have escaped the NLR. To compound the problem, when radiation forces are added, the cloud velocities are boosted even more. Yet we see clouds that are slowing down and gradually reaching systemic velocity. Therefore, the data suggest that there is a powerful force dragging on the clouds to slow their velocity.

The drag force that we introduce can have a significant effect on the clouds’ velocities. We can conclude therefore that the drag forces are a strong competitor to the radiative forces, strong enough to bring the clouds to a

halt even close to the nucleus, depending on the column densities of the outflowing clouds and the densities of the ambient medium. However the overall velocity profiles generated with radiative, gravitational, and drag forces do not match the data for NGC 1068. Assuming that the mass profile of NGC 4151 is similar to the one for NGC 1068, it will prove difficult to match its observed velocity profile, because the same linear trends are seen in the velocity of the outflowing clouds. The same can be said for Mkn 3 (Ruiz et al. 2001).

We would like to thank Douglas R. Gies and Paul J. Wiita for providing insights into some of our modeling work. We would also like to thank John. E. Everett for helpful discussions concerning the driving mechanisms of the NLR gas. The data used in this paper were obtained from the Multimission Archive at the Space Telescope Science Institute (MAST). STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. This research has made use of NASA’s Astrophysics Data System.

REFERENCES

- Antonucci, R. R. J., & Miller, J. S. 1985, *ApJ*, 297, 621
 Beckert, T., & Duschl, W. J. 2004, *A&A*, 426, 445
 Binney, J., & Merrifield, M. 1998, *Galactic Astronomy* (Princeton, NJ : Princeton University Press, 1998.)
 Bland-Hawthorn, J., Gallimore, J. F., Tacconi, L. J., Brinks, E., Baum, S. A., Antonucci, R. R. J., & Cecil, G. N. 1997, *Ap&SS*, 248, 9
 Capriotti, E., Foltz, C., & Byard, P. 1980, *ApJ*, 241, 903
 —. 1981, *ApJ*, 245, 396
 Cecil, G., Dopita, M. A., Groves, B., Wilson, A. S., Ferruit, P., Pécontal, E., & Binette, L. 2002, *ApJ*, 568, 627, [astro-ph/0112256](#)
 Chelouche, D., & Netzer, H. 2001, *MNRAS*, 326, 916
 Crenshaw, D. M., & Kraemer, S. B. 2000a, *ApJ*, 532, 247, [astro-ph/9910547](#)
 —. 2000b, *ApJ*, 532, L101, [astro-ph/0002438](#)
 Crenshaw, D. M., Kraemer, S. B., & George, I. M. 2003, *ARA&A*, 41, 117
 Crenshaw, D. M. et al. 2000, *AJ*, 120, 1731
 Das, V. et al. 2005, *AJ*, 130, 945, [astro-ph/0505103](#)
 Das, V., Crenshaw, D. M., Kraemer, S. B., & Deo, R. P. 2006, *AJ*, 132, 620
 de Vaucouleurs, G. 1948, *Annales d’Astrophysique*, 11, 247
 Dehnen, W. 1993, *MNRAS*, 265, 250
 Dopita, M. A., Groves, B. A., Sutherland, R. S., Binette, L., & Cecil, G. 2002, *ApJ*, 572, 753, [astro-ph/0203360](#)
 Dressler, A. 1984, *ApJ*, 286, 97
 Evans, I. N., Tsvetanov, Z., Kriss, G. A., Ford, H. C., Caganoff, S., & Koratkar, A. P. 1993, *ApJ*, 417, 82
 Everett, J. E. 2005, *ApJ*, 631, 689, [astro-ph/0506321](#)
 Everett, J. E., & Murray, N. 2006, [astro-ph/0610757](#)
 Ferland, G. J., Korista, K. T., Verner, D. A., Ferguson, J. W., Kingdon, J. B., & Verner, E. M. 1998, *PASP*, 110, 761
 Gallimore, J. F., Baum, S. A., & O’Dea, C. P. 2004, *ApJ*, 613, 794
 Greenhill, L. J., & Gwinn, C. R. 1997, *Ap&SS*, 248, 261
 Häring, N., & Rix, H.-W. 2004, *ApJ*, 604, L89
 Hernquist, L. 1990, *ApJ*, 356, 359
 Hutchings, J. B. et al. 1998, *ApJ*, 492, L115
 Jaffe, W. 1983, *MNRAS*, 202, 995
 Kaiser, M. E. et al. 2000, *ApJ*, 528, 260, [astro-ph/9906283](#)
 Kraemer, S. B., & Crenshaw, D. M. 2000, *ApJ*, 544, 763, [astro-ph/0007018](#)
 Krolik, J. H., & Vrtillek, J. M. 1984, *ApJ*, 279, 521
 Macchetto, F., Capetti, A., Sparks, W. B., Axon, D. J., & Boksenberg, A. 1994, *ApJ*, 435, L15
 Marconi, A., & Hunt, L. K. 2003, *ApJ*, 589, L21
 Miller, J. S., Goodrich, R. W., & Mathews, W. G. 1991, *ApJ*, 378, 47
 Nelson, C. H., Weistrop, D., Hutchings, J. B., Crenshaw, D. M., Gull, T. R., Kaiser, M. E., Kraemer, S. B., & Lindler, D. 2000, *ApJ*, 531, 257, [astro-ph/9910019](#)
 Ogle, P. M., Brookings, T., Canizares, C. R., Lee, J. C., & Marshall, H. L. 2003, *A&A*, 402, 849
 Ogle, P. M., Marshall, H. L., Lee, J. C., & Canizares, C. R. 2000, *ApJ*, 545, L81
 Parker, E. N. 1965, *Space Science Reviews*, 4, 666
 Pier, E. A., Antonucci, R., Hurt, T., Kriss, G., & Krolik, J. 1994, *ApJ*, 428, 124
 Ruiz, J. R., Crenshaw, D. M., Kraemer, S. B., Bower, G. A., Gull, T. R., Hutchings, J. B., Kaiser, M. E., & Weistrop, D. 2001, *AJ*, 122, 2961, [astro-ph/0108521](#)
 —. 2005, *AJ*, 129, 73, [astro-ph/0409754](#)
 Schmitt, H. R., Donley, J. L., Antonucci, R. R. J., Hutchings, J. B., & Kinney, A. L. 2003a, *ApJS*, 148, 327, [astro-ph/0307254](#)
 Schmitt, H. R., Donley, J. L., Antonucci, R. R. J., Hutchings, J. B., Kinney, A. L., & Pringle, J. E. 2003b, *ApJ*, 597, 768, [astro-ph/0307255](#)
 Schmitt, H. R., & Kinney, A. L. 1996, *ApJ*, 463, 498
 Schulz, H. 1990, *AJ*, 99, 1442
 Thatte, N., Quirrenbach, A., Genzel, R., Maiolino, R., & Tecza, M. 1997, *ApJ*, 490, 238
 Veilleux, S. 1991, *ApJ*, 369, 331
 Veilleux, S., Shopbell, P. L., & Miller, S. T. 2001, *AJ*, 121, 198, [astro-ph/0010134](#)
 Vrtillek, J. M. 1983, *Ph.D. Thesis*
 —. 1985, *ApJ*, 294, 121